



Diagnostica Stago, Inc.
Five Century Drive
Parsippany, NJ 07054
USA
1-800-222-COAG
973-631-1200
Fax: 973-631-1618
www.stago-us.com

ASSERACHROM VIIA-AT

BIBLIOGRAPHY

1. Silveira et al., *Relationships of plasma factor VIIa-antithrombin complexes to manifest and future cardiovascular disease* – Thromb Res 2012; 130 : 221-225
2. Spiezia et al., *Factor VIIa-antithrombin complexes plasma levels in cancer patients with and without thrombosis* – Thromb Res 2012 ; 129 : 818-819
3. Agersø et al., *Recombinant human factor VIIa (rFVIIa) cleared principally by antithrombin following intravenous administration in hemophilia patients* – J Thromb Haemost 2011; 9: 333-338
4. Agersø et al., *Clearance of rFVIIa and NN1731 after intravenous administration to Beagle dogs* – Eur J Pharm Sci 2011; 42: 578-583
5. Spiezia et al., *Factor VIIa-antithrombin complexes in patients with arterial and venous thrombosis* - Thromb Haemost 103, (2010) 1188-1192.
6. Negaard et al., *Hypercoagulability in patients with haematological neoplasia: No apparent initiation by tissue factor* - Thromb Haemost 99, (2008) 1040-1048.
7. Roubey et al., *Elevated plasma levels of factor VIIa-antithrombin complexes in patients with antiphospholipid antibodies* – J Thromb Haemost 2007; 5, Supplement 2: abstract P-T-431