

# The Fritsma Factor

YOUR INTERACTIVE HEMOSTASIS RESOURCE

Cheat Sheet—August 1, 2019



Dear Fritsma Factor participants, welcome to our August 2019 Cheat Sheet.

## July and August Quick Questions

Please refer to the results of our July 2019 Quick Question, “How does your institution manage trauma-induced coagulopathy?” For the results and discussion, tap or click [July 2019 QQ: Managing Trauma](#).

Our current Quick Question submitted by Dave McGlasson asks,

A 60-YO man taking 20 mg/day rivaroxaban for AFIB needs a high bleeding risk surgery. His creatinine is 0.9 mg/dL When should the DOAC be discontinued?

1. Never--continue treatment
2. 24 h before surgery
3. 36 h before surgery
4. 48 h before surgery
5. 60 h before surgery
6. 72 h before surgery

Please tap or click the [Fritsma Factor Home Page](#) and provide your answer.

## FF Entries

We've had a number of provocative discussions in July, including some challenging cases. Be sure to follow our Cheat Sheet links to this month's posts.

## Precision BioLogic Resources

At your convenience take a look at the list of PBI publications by navigating to the resources page or by tapping or clicking [PBI Pubs](#).

You receive our Cheat Sheet monthly. You are invited to use the site portals or contact George at [george@fritsmafactor.com](mailto:george@fritsmafactor.com) to raise new discussions or to respond to current, recent, or archived posts. If you prefer to avoid “push notifications,” contact George at the same address to request removal from our database.

Thank you for your loyalty to Fritsma Factor, thanks to Precision BioLogic for their support, and we look forward to hearing from you.

George A. Fritsma MS, MLS  
The Fritsma Factor, Your Interactive Hemostasis Resource  
[www.fritsmafactor.com](http://www.fritsmafactor.com), [george@fritsmafactor.com](mailto:george@fritsmafactor.com)